Immunization Timing

Age 2 months	
DTaP	
HepB*	
Hib	
PCV (Pneumo)	
Polio (IPV)	

Age 4 months	Interval from previous dose
DTaP	1-2 months
HepB*	1-2 months
Hib	1-2 months
PCV (Pneumo)	1-2 months
Polio (IPV)	1-2 months

Age 6 months	Interval from previous dose
DTaP	1-2 months
HepB*	2-12 months and at least 4 months after first dose
Hib	1-2 months
PCV (Pneumo)	1-2 months
Polio (IPV)	2-14 months

Age 12 months	Interval from previous dose
MMR	On or after first birthday
Varicella	On or after first birthday
Hib	2-8 months and on or after first birthday
PCV (Pneumo)	6-8 weeks and on or after first birthday

Age 15 months	Interval from previous dose
DTaP	6-12 months

Age 2 years	Interval from previous dose
НерА	2nd dose 6-12 months after first

School Recommendations Before Kindergarten

Polio, DTaP, and MMR #2. Also, HepB series and Varicella, unless already given.

Before 7th Grade

Meningitis vaccine and Tdap booster.

6-23 months and high-risk kids

Every Fall: Flu Vaccine



CALIFORNIA KIDS Love them. Immunize them.

This is a suggested schedule. For alternatives and details, consult the latest "Recommended Childhood Immunization Schedule, United States."

- * HepB series also can be started at birth, the second dose given at 1–2 months, and the third dose at 6–12 months. If using Comvax[®] (combined Hib and HepB vaccine), the third dose should be given at 12 months, not 6 months.
- † This dose can be skipped if Pedvax HIB $^\circledR$ and/or Comvax $^\circledR$ (combined Hib and HepB vaccine) are used exclusively for the infant doses.